arXiv:nucl-th/9902061v1 23 Feb 1999

Heavy-Ion Collisions and Black Holes in Anti-de-Sitter Space

John Ellis*

*Theoretical Physics Division, CERN
CH – 1211 Geneva 23*

*Invited talk given at
'RHIC Physics and beyond - Kay Kay Gee Day'
Brookhaven National Laboratory, October 23rd, 1998*

CERN-TH/98-391 nucl-th/9902061

Abstract. Recent developments linking non-perturbative quantum gauge theories in Minkowski space to classical gravity theories in anti-de-Sitter space are reviewed at a simple level. It is suggested how these spectacular advances may be extended to discuss the quark-gluon phase transition in terms of black holes in anti-de-Sitter space, with possible relevance to heavy-ion collisions.

INTRODUCTION

`Klaus Geiger was a totally special guy, and it was a privilege to have him as a friend as well as a valued collaborator. Many of us at CERN have been deeply distressed by his passing away - including secretaries, barmen and playgoers as well as physicists.`

In addition to our joint projects, Klaus and I used to enjoy shooting the bull about our other nutty ideas, not to mention string theory. I am sure he would enjoy hearing how the latest developments in this formal field might connect with his passion for RHIC physics. Mapping QCD onto gravity in anti-de-Sitter space may provide a new way of thinking about (even calculating?) the quark-hadron transition. On the way to describing this possibility in simple terms, I start by reviewing some ideas about modelling this transition that Klaus and I incorporated in our work together, and we shall see later how this may be related to anti-de-Sitter black holes.

EFFECTIVE FIELD THEORY AND THE QUARK-HADRON TRANSITION

The low-mass hadronic degrees of freedom, $\pi, K, \eta, ...$ and their low-energy interactions are described by an effective chiral Lagrangian [1], that can be written in the form

$$\mathcal{L}_{\rm eff} = -\frac{1}{2}\partial_\mu \underline{\pi} \cdot \partial^\mu \underline{\pi} + \mathcal{O}(\underline{\pi}^2(\partial\underline{\pi}^2)) + ... \quad . \tag{1}$$

The quartic and higher-order terms may in principle be calculated by integrating out quarks, or be related to the exchanges of more massive hadrons, such as the vector mesons, and their form is being constrained by data [2]. The effective Lagrangian (1) also has soliton (lump) solutions that have $I = J = 1/2$ and $B = 1$, that may be identified with baryons [3]. This picture may provide useful insight into the puzzle of the proton spin, which is purely orbital angular momentum in this soliton approach, so that the quark spins make no net contribution in the limit of light quark masses [4], in qualitative agreement with the data [5].

At finite temperatures, the chiral order parameter $\langle 0|\bar{q}q|0\rangle \to 0$ and chiral symmetry is restored [6,7]. In this limit, the chiral soliton expands: $R \to \infty$ and its mass vanishes: $M \to 0$. We therefore interpret this chiral transition also as the deconfinement transition for quarks [8]. If only the pseudoscalar degrees of freedom in (1) are taken into account, this chiral transition is second order [6,7]. However, the transition occurs at a temperature $T = 0(\Lambda_{QCD})$, so other hadronic degrees of freedom should also be taken into account [8]. Figure 1 shows how an order parameter χ related to the gluon condensate $\langle 0|G_{\mu\nu}G^{\mu\nu}|0\rangle$ behaves as a function of temperature when these are included. At low temperature $T_0 < T_1$, the free energy is minimized when $\chi \neq 0$ and the heavier hadrons with $m \propto \chi$ are massive. At T_1, the free energy is equal in this and the $\chi = 0$ phase where the hadrons are all massless. Below T_2, there is the possibility of a mixed phase, depending on the history of the system. Finally, when one heats beyond T_2, the $\chi \neq 0$ confined vacuum ceases to exist, and the theory must be in the unconfined phase. This picture is reminiscent of a liquid-gas transition [9], and is analogous with phase transitions of gravity in anti-de-Sitter space, as we shall see later.

Klaus and I incorporated these ideas [10] in his space-time parton-shower Monte Carlo [11], to provide an afterburner for hadronization. To see how this works, consider the process $e^+e^- \to Z^0 \to \bar{q}q$, where the initial decay produces a pair of "hot"



FIGURE 1. *Qualitative variation of the free energy with temperature in QCD [8]. At temperatures below T_0, there is no stable state at the origin, the only stable state (with non-zero order parameter) corresponds to confinement, and there is a string tension. There is no stable confined state at temperatures above T_2. Between T_0 and T_2, there is the possibility of a mixed phase. The confined and unconfined phases have equal free energies at the temperature T_1, which would be T_c under adiabatic conditions.*

off-shell partons as shown in Fig. 2a. These create a local "hot spot" of the unconfined phase, which is surrounded by the usual "cold" hadronic vacuum. During the subsequent parton-shower development, the "hotter" deeply-virtual partons decay into "cooler" partons closer to mass shell, as the "hot spot" expands and cools. Since an isolated parton is impossible, whenever a parton becomes $\gtrsim 1$ fm away from its nearest neighbour, it should be confined and hadronize. This process continues until all the partons "cool", separate and hadronize, as seen in Fig. 2a for $e^+e^- \to Z^0 \to \bar{q}q$ and in Fig. 2b for $e^+e^- \to W^+W^- \to \bar{q}q\bar{q}q$.

We modelled [10] the parton-hadron conversion using the chiral Lagrangian (1) to describe the "cool" phase, and describing the transition by analogy with the finite-temperature transition shown in Fig. 1, but replacing temperature T by the separation L between a parton pair: $T \leftrightarrow \frac{1}{L}$ [12]. We interpreted L as the effective size of the local cooling region, and treated the hadronic degrees of freedom as if they were in a box of this size. We then estimated heuristically the hadronization probability $P(L)$ by calculating the tunnelling through the barrier shown in Fig. 1 between the "partonic" $\chi = 0$ and hadronic $\chi \neq 0$ vacua. This probability peaks at a characteristic separation $L \sim 0.6$ to 0.8 fm. We have applied this picture


$\bar{q}$
q
91.5 GeV
$e^+e^- \longrightarrow Z^0 \longrightarrow q\bar{q}$
$\bar{q}'$
$\bar{q}$
183 GeV
q
q'
$e^+e^- \longrightarrow W^+W^- \longrightarrow q\bar{q}'\ q'\bar{q}$

FIGURE 2. *Schematics of the two e^+e^- event types $e^+e^- \to Z^0 \to \bar{q}q$ and $e^+e^- \to W^+W^- \to \bar{q}q\bar{q}q$. The final-state hadron distribution in Z^0 events (left) is due to exclusively 'endogamous' hadronization of the partonic offspring from the $q\bar{q}$ dijet [10], whereas in W^+W^- events (right) there is, in addition, the possibility of 'exogamous' hadron production involving a mating of partons from the two different $W^+ \to q\bar{q}$ and $W^- \to q\bar{q}$ dijets [14].*

to various physical conditions, including deep-inelastic scattering [13] and $e^+e^- \to W^+W^- \to$ hadrons [14], as well as to $e^+e^- \to Z^0 \to$ hadrons [10]. An afterburner for Bose-Einstein correlations [15] that we explored together with Klaus is described here by Ulrich Heinz and Urs Wiedemann [16].

In the rest of this talk, we explore a different direction, first looking at more theoretical approaches to confinement, and seeing later their possible relation to the ideas discussed in this section.

CONFINEMENT AND MONOPOLES

A formal approach to confinement is based on the space-like Wilson loop integral

$$W(C) = e^{ie\int_c A\cdot dl} \ , \tag{2}$$

and the time-like Polyakov loop integral

$$P(C') = e^{ie\int_{C'} A\cdot dl} \ . \tag{3}$$

If we consider a curve C' that has the form of a rectangle with time-like sides of duration t separated by a distance R, in the confining phase we expect that

$$P(R,t) \sim e^{-\mu R t} \ , \tag{4}$$

corresponding to an effective potential

$$V(R) \sim \mu R \tag{5}$$

This is believed to correspond to a flux tube in which the chromoelectric flux lines have been squeezed by a dual analogue of the Meissner effect in conventional superconductivity [18]. In the conventional case, there is a condensate of Cooper pairs $\langle e^- e^- \rangle$ that seeks to expel magnetic flux lines. In the QCD case, there should be a condensate of chromomagnetic monopoles that seek to expel and squeeze chromoelectric flux. Essential aspects of this picture have been demonstrated in $N = 2$ supersymmetric gauge theory, which has an exact non-perturbative solution [17].

The chromomagnetic monopoles arise in the projection of a non-Abelian gauge theory onto its maximal Abelian subgroup, e.g., $SU(N) \to U(1)^{N-1}$ [18]. It is believed that this Abelian subgroup may be solely responsible for confinement, and Polyakov has proposed a formal string description of QCD in this confined phase [19]. Assuming monopole condensation at a characteristic QCD scale Λ, he has argued that low-energy physics is described by an effective action for a chiral gauge field, that includes an antisymmetric tensor field coupled to string:

$$A \to \int d^4x B_{\mu\nu}(x) \int d^2\sigma \epsilon_{ab} \partial_a X^\mu \partial_b X^\nu \delta^4(X(\sigma) - x) \tag{6}$$

where σ and the suffices a, b denote world-sheet coordinates. Relating this to the formalism of fundamental string theory, and deriving a full theory of QCD string, remain theoretical challenges.

INTRODUCTION TO STRING THEORY

It is worth recalling that string models were born from old-fashioned strong-interaction theory. The lowest-lying hadrons p, n, π were found to be accompanied by many heavier hadronic states, whose maximum spins grew quadratically with their masses:

$$m^2 \propto J \ , \tag{7}$$

as would occur in a linearly-rising potential (5). Such a potential would lead to an infinite set of unstable excited states.

It was observed that such an infinite set of states would make possible a dual bootstrap picture in which the infinite set of direct-channel resonances could be resummed to yield an infinite set of cross-channel exchanges, which were associated with Regge trajectories. A mathematical model scattering amplitude embodying these ideas was found by Veneziano [20].

Shortly afterwards, it was shown that this amplitude could be derived from a string theory [21], in which the highest-spin excited states had

$$J \simeq \alpha' m^2 \ , \tag{8}$$

where $\alpha' \sim (1 \text{ GeV})^{-2}$ is the string tension. The hadrons could be interpreted in this picture as excited vibrational states of open strings with massless quarks attached to their ends. Unitarity at higher orders required that these be supplemented by closed strings that were thought to be related to the Pomeron and to glueballs.

Attention to this approach to the strong interactions was distracted by its failure to explain the point-like parton structures that had emerged in high momentum-transfer processes. Interest in the string approach to QCD lapsed with the discovery of QCD, which showed how this point-like behaviour could be understood in field theory through the property of asymptotic freedom.

However, around the same time it was proposed that string theory be reinterpreted as a quantum theory of gravity and all the other particle interactions [22]. This proposal arose from the observation that the spectrum of closed string included a massless spin-2 particle that could be interpreted as the graviton, as well as massless spin-1 particles that could be interpreted as gauge bosons. This reinterpretation apparently required rescaling the string tension: $\alpha' \to \mathcal{O}\left(\frac{1}{m_P^2}\right)$, where $m_P \simeq 10^{19}$ GeV is the energy scale at which four-dimensional gravitational interactions become strong.

String theories can be written in terms of two-dimensional field theories on the world sheet which describes the propagation of the string through time. A scalar field on the world sheet of a closed string can be decomposed into left- and right-moving waves:

$$\phi(r+t) + \phi(r-t) \ , \tag{9}$$

each of which can be expanded in normal modes. Such a two-dimensional field theory is relatively easy to make finite at the loop level, thus offering the prospect of a finite quantum theory of gravity. This finiteness may be traced to the softening of divergence difficulties that occurs when point particles are replaced by extended structures, e.g.:

$$\int d^4k \frac{1}{k^2} \leftrightarrow \int_{1/\Lambda = R} d^4x \frac{1}{x^6} \sim \frac{1}{R^2} \ , \tag{10}$$

where R is the size of the extended object. Strings are the simplest examples of this approach, but why not also consider higher-dimensional internal structures? This is actually what happens when one confronts non-perturbative aspects of string theory [23], as we discuss later.

In making a finite string theory, however, one must be careful to avoid anomalies at the quantum level. In the case of a purely bosonic string with no additional world-sheet degrees of freedom, at the perturbative level this requires embedding it in 26 space-time dimensions [24]. If the string is made supersymmetric, the necessary number of dimensions is reduced to ten. The complete quantum consistency of these theories was demonstrated by Green and Schwarz [25], triggering the first string revolution.

The original bosonic string suffered from the defects that it had no fermionic matter and that its vacuum was unstable, since it had a tachyonic scalar field. Superstrings gained on both counts, since they contained fermions and had stable vacua with zero cosmological constant. However, they were initially discarded because it was not seen how to incorporate parity violation. This was first achieved with the ten-dimensional heterotic string [26], which retained the feature of a stable zero-energy vacuum. The final step towards making realistic models came with the realization that one could construct four-dimensional heterotic string models, either by compactifying the six surplus dimensions on a small manifold with characteristic size $\sim \frac{1}{m_P}$ [27], or by direct construction in four dimensions [28]. In the latter case, one replaced the surplus dimensions by internal world-sheet degrees of freedom. Such models could accommodate a realistic GUT-like or Standard-Model-like gauge group in four dimensions [29].

Perturbative string models had a number of phenomenological successes. They predicted that there should be at most ten dimensions, in agreement with the observed number of four. They predicted that the gauge group should have rank no greater than 22, in agreement with the rank four of the Standard Model. They could not accommodate matter in large representations of the Standard Model gauge group, such as the $\underline{3}$ of SU(2) or the $\underline{8}$ of SU(3), also in agreement with experiment. Also, in a generic string model the maximum size of a Yukawa coupling λ, such as that of the top quark, is of the same order as the gauge coupling g [30]. One therefore predicts a maximum quark mass $m_t = \mathcal{O}(g)\langle H \rangle$, which yields the successful prediction $m_t \lesssim 190$ GeV after including renormalization-group effects.

Perturbative string theories also permit a calculation of the gauge unification scale from first principles [31]:

$$m_{\rm GUT} = 0(g) \times \frac{m_P}{\sqrt{8\pi}} \simeq \text{few} \times 10^{17} \text{ GeV} . \qquad (11)$$

This top-down calculation is close to the bottom-up estimate of $m_{\rm GUT}$ based on the experimental values of the gauge couplings: $m_{\rm GUT} \sim 2 \times 10^{16}$ GeV [32]. but there is an apparent discrepancy by over an order of magnitude. This provided motivation for considering strongly-coupled string models with an extra (relatively) large spatial dimension in the context of M theory [33], as we discuss in the next section.

BEYOND STRINGS

The second string revolution was triggered by the understanding of non-perturbative effects in string theory [23], and the role of higher-dimensional extended objects such as membranes, which do not appear in perturbative string theory. The lowest two-dimensional membrane describes a three-dimensional world volume as it propagates, and there are analogous higher-dimensional extended structures. These may be obtained as soliton solutions of string theory, with masses $m \sim 1/g_s$. Therefore, they become light in the strong-coupling limit, just as magnetic monopoles become light in the strong-coupling limit of gauge theory. It has been known for some time that there is a duality in gauge theory between these monopoles and conventional gauge particles [34]. Now it is known that similar dualities exist in string theory: solitons in some strongly-coupled theories can be identified with light states in the weakly-coupled perturbative limit of some other string theory [35]. A related development has been the realization that the string coupling g_s should be interpreted as the expectation value of a quantum field, that can be interpreted as an extra dimension. In the strong-coupling limit $g_s \to \infty$, this extra dimension also becomes large: $R \to \infty$, and a ten-dimensional perturbative string theory becomes a non-perturbative eleven-dimensional theory [33]. It is now believed that all string theories are related by dualities, and that they all may be regarded as particular limits of this underlying eleven-dimensional theory, known as M theory. In the low-energy limit, this theory reduces to eleven-dimensional supergravity [36], but its full spectrum is infinite.

The question now arises: which limit of M theory best describes the physics we know? An important clue is provided by the bottom-up calculation of the unification scale using the low-energy gauge-coupling measurements at LEP and elsewhere [32]. This can be reconciled with the top-down string calculation [31] if the eleventh dimension is relatively large: $L_{11} \gg 1/m_{\mathrm{GUT}} \sim 1/(10^{16}\ \mathrm{GeV}) \gg 1/m_P$. The idea is that the gravitational interactions "feel" this extra dimension, and so become stronger at shorter distances: the Newton potential $V_N \propto 1/r \to 1/r^2$ in five dimensions. In this way, the gravitational coupling may rise more quickly at energies $E \gtrsim 1/L_{11}$, so as to be unified with the gauge interactions (that do not feel the extra dimension) as shown in Fig. 3. The proposal is that the gauge interactions act only on ten-dimensional subspaces of the full eleven dimensions, that may be regarded as "capacitor plates" separated by the large eleventh dimension in which gravity acts, as seen in Fig. 4 [33]. Six dimensions are compactified, as in weakly-coupled string theory, with a size $R \sim 1/m_{\mathrm{GUT}}$. Thus physics is described by a five-dimensional effective supergravity theory at energies



FIGURE 3. *Sketch of the possible evolution of the gauge couplings* α_i *and the gravitational coupling* G*: if there is a large fifth dimension with size* $\gg m_{GUT}^{-1}$*,* G *may be unified with the gauge couplings at the GUT scale.*

$L_{11}^{-1} < E < m_{\rm GUT}$ [37].

AN EXTRA DIMENSION FOR GAUGE THEORIES?

We saw in the previous section how a ten-dimensional string theory becomes effectively an eleven-dimensional theory in the strong-coupling limit. Does something similar also happen in gauge theories [38]? Such a possibility would be consistent with the holographic principle [39], according to which the information in a field theory in some space-time can be encoded in another field theory on its boundary. This was proposed originally in the context of quantum gravity, as a way of understanding the fact that the black-hole entropy is proportional to the area of its event horizon. In other words, all the information in the bulk of a black hole is encoded on its surface. In the present context, information about the quantum behaviour of a gauge theory is conjectured to be encoded in the classical behaviour of a higher-dimensional gravity theory [38,40].

The crucial step is to recall that Minkowski space in d dimensions M_d is itself the boundary of anti-de-Sitter (AdS) space in $d+1$ dimensions ($\mathrm{AdS_{d+1}}$), which is described by the metric



FIGURE 4. *The capacitor-plate scenario favoured in eleven-dimensional M theory. The eleventh dimension has a size* $L_{11} \gg M_{GUT}^{-1}$, *whereas dimensions* $5, ..., 10$ *are compactified on a small manifold K with characteristic size* $\sim M_{GUT}^{-1}$. *The remaining four dimensions form (approximately) a flat Minkowski space* M_4.

$$ds^2 = \left(1 + \frac{r^2}{b^2}\right) dt^2 + \frac{1}{(1 + \frac{r^2}{b^2})} dr^2 + r^2 d\Omega^2 \ , \tag{12}$$

where b is the AdS radius. Is gauge theory on M_d equivalent to some theory on AdS$_{d+1}$? This bulk theory is likely to be a gravity theory, because of the Kaluza-Klein trick:

$$G_{MN}(M, N = 0, 1, 2, 3, 4) \to g_{\mu\nu}, g_{\mu 4}, g_{4\nu}, g_{44} \ , \tag{13}$$

according to which the $(\mu 4, 4\nu)$ components of g_{MN} in five dimensions are identified as four-dimensional gauge fields.

Thus we arrive at the Maldacena conjecture [38]: that a d-dimensional conformal (i.e., finite, with no divergences) supersymmetric gauge theory may be equivalent, at least in a certain limit, to a conformal supergravity (or string) theory in $(d+1)$-dimensional AdS space. The appropriate limit in which the conjecture was first formulated was that of large N_c for the gauge theory, where the AdS$_{\rm d+1}$ curvature becomes small and reliable classical solutions to the supergravity field equations can be found. The spectrum of supergravity excitations in ADS$_{\rm d+1}$ was found to correspond to the spectrum of an $\mathcal{N} = 4$ supersymmetric gauge theory in the large-

N_c limit, which is known to be finite. It was also suggested that a similar analysis could be made for non-conformal field theories.

Witten extended this idea to the $\mathcal{N} = 4$ supersymmetric theory at finite temperature $t \neq 0$ [41], and was able to demonstrate magnetic confinement as discussed in Section 3, with the appearance of a mass gap. Since the boundary conditions used to study finite-temperature field theory violate supersymmetry, this provides a prototype for studying four-dimensional non-supersymmetric gauge theories at large N_c. However, it was not possible in [41] to demonstrate asymptotic freedom, and hence that a different phase did not appear in the weak-coupling limit.

A NON-CONFORMAL(IST) APPROACH

Conformal symmetry is very powerful: with no infinities and hence no need to introduce a renormalization scale, one can derive many rigorous results. However, the real world is not described by a conformal gauge theory: there is asymptotic freedom, there are quark and gluon condensates, etc.. So how can one extend the above ideas to go beyond conformal symmetry [42,43]?

In string theory, the conformal factor ϕ in the world-sheet metric

$$\gamma_{ab} = e^{\phi}\hat{\gamma}_{ab}, \tag{14}$$

where $\hat{\gamma}_{ab}$ is a fiducial reference metric, is called a Liouville field [44]. Perturbatively it decouples in a conformal string theory such as the bosonic string in 26 dimensions or the superstring in ten dimensions, but has non-trivial dynamics in a non-conformal string theory, as would be needed to describe QCD, which lives in fewer dimensions. We have suggested that one identifies the Liouville field with a q-number scale factor, that can be identified as a dynamical renormalization scale in non-critical string theory [45]. We have also discussed the interpretation of world-sheet defects [46] in such a non-critical Liouville string theory as black holes in target space [47].

These ideas have recently been applied to the problem of confinement in gauge theory [43]. We have shown that a world-sheet vortex defect corresponds to the intersection of a target-space magnetic-field line with the world sheet, as seen in Fig. 5. This is because one may rewrite the line integral in (2) in the following way:

$$\int_C A \cdot dl = \int_{\Sigma(C)} B \cdot ds \ , \tag{15}$$

where dS is an element of the surface $\Sigma(c)$ that is bounded by the curve C whose line element is dl. One may then decompose

$$B = \epsilon_{ab}\partial_a A_b = \epsilon_{ab}\partial_a\partial_b X^\mu A_\mu(x) + \dots , \tag{16}$$

where $\partial_a\partial_b X^\mu \neq 0$ in the presence of a world-sheet vortex. Condensation of magnetic monopoles in target space creates a flux through the world sheet that corresponds to condensation of vortices on the world sheet. This condensation is known to occur at a low enough temperature $T < T_{\mathrm{vortex}}$, and we have shown that vortex condensation generates a tension for the string:

$$\mu = (\alpha')^{-1} \propto \langle (\epsilon_{ab}\partial_a\partial_b X)^2 \rangle \neq 0 \tag{17}$$

corresponding to confinement in the string picture [43].



FIGURE 5. *A string world sheet $\Sigma(C)$, whose boundary is a Wilson loop C, in the presence of a four-dimensional space-time magnetic field, as could be generated by a target-space monopole. The intersection of the field line with the string world sheet results in a world sheet vortex, which is related by world-sheet duality to a world-sheet spike. Condensation of magnetic flux in space time corresponds to condensation of vortices on the world sheet, which generates a string tension [43].*

We have also shown [43] how these ideas can be extended to AdS black holes, as illustrated in Fig. 6. Black holes are related to defects of different type, called spikes, that are related to world-sheet vortex defects by a world-sheet duality [46]. Stable AdS black holes correspond to a condensation of spikes on the world sheet, which is known to occur at a high enough temperature $T > T_{\mathrm{spike}} > T_{\mathrm{vortex}}$. The AdS gravity theory has phases [48] that parallel those of QCD:

$$T < T_{\mathrm{vortex}} : \qquad \text{vortices condense, no black holes, confinement}$$

$$\begin{array}{rl} T_{\rm vortex} < T < T_{\rm spike}: & \text{vortices unbound, black holes unstable, mixed phase} \\ T_{\rm spike} < T: & \text{spikes condense, stable black holes, deconfinement} \end{array} \tag{18}$$

The temperature dependence of the free energy resembles that shown in Fig. 1 for QCD, with $T_2 \leftrightarrow T_{\rm spike}$ and $T_0 \leftrightarrow T_{\rm vortex}$. The behaviour of the bulk theory is reminiscent of a liquid-gas transition [9], a possibility to which we return shortly.



FIGURE 6. *The string world sheet of Fig. 5 is embedded in four-dimensional Minkowski space M_4, which is regarded here as the boundary of five-dimensional Anti-de-Sitter space AdS_5. A black hole in the bulk of AdS_5 corresponds to a vortex on the world sheet, whose condensation corresponds to confinement [43].*

WHAT DOES THIS HAVE TO DO WITH HEAVY IONS?

These advanced new theoretical ideas certainly provide deeper understanding and insight into confinement and deconfinement in QCD [40]. They may also provide useful new calculational techniques. Concretely, one may hope to tackle non-perturbative strong-coupling phenomena in four-dimensional QCD using the techniques of classical weakly-coupled gravity or string theory in AdS_5. Indeed, people have started to calculate quantities of experimental interest.

Glueball Masses [49]: correlation functions in QCD_4 in the limit of large N_c can be related to classical properties of supergravity in AdS_5. Particle poles can therefore be related to classical eigenvalue equations. In this way, calculations in a certain limit have been made of the ratios of glueball masses, e.g.,

$$\frac{m_{0^{-+}}}{m_{0^{++}}} = 1.20 \ , \tag{19}$$

to be compared with the ratio 1.36 ± 0.32 obtained from a lattice computation. It remains to be seen how results in the AdS_5 supergravity limit correspond to the physical case of QCD_4, but at least one has some numbers to discuss.

Vacuum Properties [50]: First calculations have been made of the gluon condensate and topological susceptibility χ_t, with the results:

$$\frac{\langle 0|GG|0\rangle}{\langle 0|GG|0\rangle|_{\mathrm{lattice}}} \simeq 0.9 \ , \quad \frac{\chi_t}{\chi_t|_{\mathrm{lattice}}} \simeq 1.7 \ , \tag{20}$$

as well as the string tension:

$$\frac{\mu}{\mu|_{\mathrm{lattice}}} \simeq 0.4 \ . \tag{21}$$

These numbers are not perfect, and the AdS_5 supergravity limit does not correspond to the physical QCD_4 limit, but please remember that these are early days yet.

Heavy-Quark Potential [51]: First calculations are quite promising, with the expected form:

$$V(R) \sim \mu R + (\mathrm{cont.}) + \exp(-R\Lambda_{QCD}) + \ldots \ , \tag{22}$$

emerging.

Finally, what about heavy ions? We are trying to model the behaviour of the free energy and pressure near the QCD transition [9], which are known from lattice calculations to exhibit departures from an ideal gas. This effect is particularly marked for the pressure, which is considerably less than the ideal-gas value, as seen in Fig. 7 [52]. One interpretation of this departure has been in terms of an effective magnetic mass for the gluon above the transition temperature.

We regard this behaviour as a manifestation of a liquid-gas transition for black holes in AdS_5, which are regarded as effective regulators for the monopoles in M_4. These black holes interact via a non-trivial effective potential with analogues of van der Waals forces. A standard analysis then leads to a van der Waals formula for the equation of state,

$$RT = (V - b)\left(P + \frac{a}{v^2}\right) \ , \tag{23}$$

leading to the familiar pattern of isothermal pressure curves. Based on this picture, we obtain the behaviours of the free energy and pressure shown in Fig. 8 near the



FIGURE 7. *Scaled energy density ϵ/T^4 compared with three times the scaled pressure $3p/T^4$, as obtained from lattice QCD calculations: for a discussion and original references, see [52].*

critical temperature T_c [9]. These have the right qualitative behaviour, including the vanishing of the pressure close to T_c.

We conclude on the optimistic note that modern ideas [23,38,43] in non-perturbative string theory duality and conformal field theory may indeed have some relevance to heavy-ion physics. Some non-perturbative QCD quantities have already been estimated [49–51], and more are on the way. Not only qualitative insights but also quantitative results may eventually emerge. The finite-temperature QCD transition may be yielding to the theoretical assault [9]. Perhaps these ideas may even eventually enable the calculation of physical quantities measurable directly in heavy-ion collisions?

REFERENCES

1. J. Gasser and H. Leutwyler, *Phys. Rep.* 87 (1987) 77.
2. A. Pich, hep-ph/9806303.
3. E. Witten *Nucl.Phys.* B223 (1983) 422, 433;
   G.S. Adkins, C. Nappi and E. Witten, *Nucl.Phys.* B228 (1983) 552.
4. S.J. Brodsky, J. Ellis and M. Karliner, *Phys.Lett.* B206 (1988) 309.
5. Spin Muon Collaboration, B. Adeva et al., *Phys.Rev.* D58 (1998) 112002.
6. P. Binetruy and M.K. Gaillard, *Phys.Rev.* D32 (1985) 931.



FIGURE 8. *Scaled energy density and three times the scaled pressure obtained from a preliminary analysis using an effective van der Waals description of a non-ideal gas of* AdS_5 *black holes [9]. The qualitative behaviour, in particular the fact that the scaled energy density is larger than* $3p/T^4$ *in the transition region, is similar to that in the lattice calculations shown in Fig. 7.*

7. J. Gasser and H. Leutwyler, *Phys.Lett.* 184B (1987) 83 and 188B (1987) 477.
8. B.A. Campbell, J. Ellis and K.A. Olive, *Phys.Lett.* B235 (1990) 325 and *Nucl.Phys.* B345 (1990) 57.
9. J. Ellis, A. Ghosh and N.E. Mavromatos, CERN preprint TH/99-43 in preparation.
10. J. Ellis and K. Geiger, *Nucl.Phys.* A590 (1995) 609c and *Phys.Rev.* D52 (1995) 1500.
11. K. Geiger, *Comput.Phys.Commun.* 104 (1997) 70.
12. J. Gasser and H. Leutwyler, *Nucl.Phys.* B307 (1988) 763.
13. J. Ellis, K. Geiger and H. Kowalski, *Phys.Rev.* D54 (1996) 5443.
14. J. Ellis and K. Geiger, *Phys.Rev.* D54 (1996) 1967 and *Phys.Lett.* B404 (1997) 230.
15. K. Geiger, J. Ellis, U. Heinz and U. Wiedemann, hep-ph/9811270.
16. U. Heinz and U. Wiedemann, talks at this meeting.
17. N. Seiberg and E. Witten, *Nucl.Phys.* B431 (1994) 484.
18. S. Mandelstam, *Phys.Rep.* 23C (1976) 245;
    G. ‘t Hooft, *Nucl.Phys.* B190 [FS3] (1981) 455.
19. A.M. Polyakov, *Nucl.Phys.* B72 (1974) 461; *Phys.Lett.* B59 (1975) 79; *ibid.* B268 (1986) 406; *Gauge Fields and Strings* (Harwood, 1987); hep-th/9711002.
20. G. Veneziano, *Nuovo Cim.* A57 (1968) 190.

21. P. Goddard, J. Goldstone, C. Rebbi and C. Thorne, *Nucl.Phys.* B56 (1973) 109.
22. J. Scherk and J.H. Schwarz, *Nucl.Phys.* B81 (1974) 118.
23. J. Polchinski, *Phys.Rev.Lett.* 75 (1995) 184;
    J. Polchinski, S. Chaudhuri and C. Johnson, hep-th/9602052 and references therein;
    J. Polchinski, TASI lectures on D-branes, hep-th/9611050, and references therein.
24. C. Lovelace, *Phys.Lett.* 34B (1971) 500.
25. M.B. Green and J.H. Schwarz, *Phys.Lett.* 149B (1984) 117 and *Phys.Lett.* 151B (1985) 21.
26. D.J. Gross, J.A. Harvey, E. Martinec and R. Rohm, *Phys.Rev.Lett.* 54 (1985) 502.
27. P. Candelas, G. T. Horowitz, A. Strominger and E. Witten, *Nucl.Phys.* B258 (1985) 46.
28. See, e.g., I. Antoniadis, C. Bachas and C. Kounnas, *Nucl.Phys.* B289 (1988) 87;
    I. Antoniadis and C. Bachas, *Nucl.Phys.* B298 (1987) 586.
29. See, e.g., I. Antoniadis, J. Ellis, J.S. Hagelin and D.V. Nanopoulos, *Phys.Lett.* B231 (1989) 65.
30. E. Witten, *Phys.Lett.* B155 (1985) 151.
31. See, e.g., I. Antoniadis, J. Ellis, R. Lacaze and D.V. Nanopoulos, *Phys.Lett.* B268 (1991) 188.
32. J. Ellis, S. Kelley and D.V. Nanopoulos, *Phys.Lett.* B249 (1990) 441 and *Phys.Lett.* B260 (1991) 131;
    U. Amaldi, W. de Boer and H. Furstenau, *Phys.Lett.* B260 (1991) 447;
    P. Langacker and M. Luo, *Phys.Rev.* D44 (1991) 817.
33. P. Horava and E. Witten, *Nucl.Phys.* B460 (1996) 506;
    E. Witten, *Nucl.Phys.* B471 (1996) 135;
    P. Horava and E. Witten, *Nucl.Phys.* B475 (1996) 94;
    T. Banks and M. Dine, *Nucl.Phys.* B479 (1996) 173.
34. C. Montonen and D.I. Olive, *Phys.Lett.* 72B (1997) 117.
35. For a review, see: A. Sen, hep-ph/9810356.
36. E. Cremmer, B. Julia and J. Scherk, *Phys.Lett.* 76B (1978) 409;
    E. Cremmer and B. Julia, *Phys.Lett.* 80B (1978) 48 and *Nucl.Phys.* B159 (1979) 141.
37. I. Antoniadis, S. Ferrara and T. Taylor, *Nucl.Phys.* B460 (1996) 489;
    A. Lukas, B.A. Ovrut and D. Waldram, hep-th/9803235;
    A. Lukas, B.A. Ovrut, K.S. Stelle and D. Waldram, hep-th/9806051;
    R. Donagi, A. Lukas, B.A. Ovrut and D. Waldram, hep-th/9811168;
    J. Ellis, Z. Lalak, S. Pokorski and W. Pokorski, *Nucl.Phys.* B540 (1999) 149;
    J. Ellis, Z. Lalak and W. Pokorski, hep-th/9811133.
38. J. Maldacena, *Adv.Theor.Math.Phys.* 2 (1998) 231.
39. G. ‘t Hooft, gr-qc/9310026;
    L. Susskind, hep-th/9409089.
40. For a theoretical review of this explosive field, see:
    J.L. Petersen, hep-th/9902131.
41. E. Witten, *Adv.Theor.Math.Phys.* 2 (1998) 505.
42. S. Gubser, I. Klebanov and A.M. Polyakov, hep-th/9802109.
43. J. Ellis and N.E. Mavromatos, CERN-TH-98-264, hep-th/9808172.
44. F. David, *Mod.Phys.Lett.* A3 (1988) 1651;

J. Distler and H. Kawai, *Nucl.Phys.* B321 (1989) 509.

45. J. Ellis, N.E. Mavromatos and D.V. Nanopoulos, *Phys.Lett.* B293 (1992) 37; *Mod.Phys.Lett.* A10 (1995) 425;
    Lectures presented at the *Erice Summer School, 31st Course: From Supersymmetry to the Origin of Space-Time*, Ettore Majorana Centre, Erice, July 4-12 1993 ; hep-th/9403133, 'Subnuclear Series' Vol. 31, (World Scientific, Singapore 1994), p.1
46. J. Cardy, *Nucl.Phys.* B205 (1982) 17;
    J. Cardy and E. Rabinovici, *Nucl.Phys.* B205 (1982), 1;
    B. Ovrut and S. Thomas, *Phys.Lett.* B237 (1991) 292;
    J. Ellis, N.E. Mavromatos and D.V. Nanopoulos, *Phys.Lett.* B289 (1992) 25 and *Mod.Phys.Lett.* A12 (1997) 2813.
47. J. Ellis, N.E. Mavromatos and D.V. Nanopoulos, *Int.J.Mod.Phys.* A12 (1997) 2639.
48. S. Hawking and D. Page, *Comm.Math.Phys.* 87 (1983) 577.
49. C. Csáki, H. Ooguri, Y. Oz and J. Terning, preprint UCB-PTH-98-30, hep-th/9806021;
    R. de Mello Koch, A. Jevicki, M. Mihailescu and J. P. Nunes, preprint BROWN-HET-1130, hep-th/9806125;
    M. Zyskin, *Phys.Lett.* B439 (1998) 373;
    H. Ooguri, H. Robins and J. Tannenhauser, *Phys.Lett.* B437 (1998) 77;
    J. Minahan, hep-th/9811156.
50. A. Hashimoto and Y. Oz, hep-th/9809106.
51. J. Greensite and P. Olesen, *JHEP* 9808 (1998) 009 and hep-th/9901057.
52. For a recent review, see: H. Satz, hep-ph/9711289.